

# Urinary hCG follow-up<sup>1</sup>


**checkToP<sup>®</sup>** is an urinary test which determines the outcome of the treatment after a previously confirmed Termination of Intrauterine Pregnancy (ToP).



**checkToP<sup>®</sup> is not a classical pregnancy test.**

It detects a hCG level of 1 000 mIU/mL or more in urine.

**Medical termination of pregnancy (MToP) consists in several steps:**



- If you perform the test too early, the level of hCG rate could still be too high and lead you to think that Termination of Pregnancy failed.

- On the other way, if you perform this test too late, and in the exceptional case where the termination would have failed, you would delay the management of your care and support.

## PROCEDURE

**checkToP<sup>®</sup>** must be performed on the first morning urine that should be more concentrated.

- 1 Hold the absorbent tip directly in your urine stream or collect your urine in a clean and dry container and dip the absorbent tip into the urine for at least 5 to 10 seconds. **Take care no urine enters in the test window.**
- 2 Lay the test device on a dry flat surface with the two windows facing up.
- 3 Wait between **5 to 10 minutes** to read the test results.

**Important: Do not read before 5 minutes or after 10 minutes.**



## RESULT INTERPRETATION



One band in the control window and no band in the test window

**Termination of pregnancy is confirmed\***



One band in each reading window

**Treatment for medical termination of pregnancy may not have worked or is not complete**

**=> Consult your physician**

**(please report it to [pv@exelgyn.com](mailto:pv@exelgyn.com))**



or



No band in the control window, with or without a band in the test window: **INCONCLUSIVE test**

**(please report it to [pv@exelgyn.com](mailto:pv@exelgyn.com))**

Test to be repeated with a new **checkToP<sup>®</sup>** and fresh urine sample obtained 24 hours later after the inconclusive test

**\*The test is 99% reliable. However, even if the test confirms the termination of pregnancy, you should still contact your physician if you have the following symptoms:**



**No bleeding or limited bleeding since the last treatment intake**



**Tummy growing**



**Feeling sick**



**Tender breasts**



**No period by 1 month after treatment**

**Consult your physician to have additional examinations**